

Influence of naringenin on the activity of enzymes participating in steroidogenesis in male rats

Papież M A

Department of Cytobiology and Histochemistry, Pharmaceutical Faculty, Medical College,
The Jagiellonian University, Kraków, Poland

Abstract

The aim of the experiment was to histochemically examine the activity of the following selected steroidogenic enzymes: $\Delta^3\beta$ HSD and 17β HSD and the housekeeping enzyme G6PDH, in experimental rats after subcutaneous injections of naringenin flavonoid, at a daily dose of 15 mg/kg of body mass. The enzyme activities were measured by the microdensitometric method. Additionally, radioimmunological assay for testosterone level was conducted in homogenates of testes. A significant decrease in the activity of 17β HSD and of G6PDH was found, while the activity of $\Delta^3\beta$ HSD was not significantly changed. The results permit a statement that naringenin causes minor changes in metabolic processes in the testes of rats but it does not significantly affect the synthesis of androgens.

Key words: $\Delta^3\beta$ -hydroxysteroid dehydrogenase,
 17β -hydroxysteroid dehydrogenase,
glucose-6-phosphate dehydrogenase,
histochemistry, testis, rat, naringenin.

Introduction

Naringenin is a biphenolic compound, belonging to the group of flavonoids and to the class of flavanones. This flavonoid occurs in greatest quantities in citrus fruits, as well as in hop [1]. Because of its affinity to the estrogen receptor α and the ability to stimulate proliferation of cells in female reproductive tracts, this flavonoid is placed among the phytoestrogens,

i.e., non-steroid compounds of plant origin, showing a structural similarity with 17β -estradiol [2, 3]. Like other phytoestrogens, naringenin inhibits the activities of 17β HSD and aromatase *in vitro* [4]. The objective of this study was to examine the *in vivo* effects of naringenin on the activity of selected steroidogenic enzymes: $\Delta^3\beta$ HSD and 17β HSD, as well as G6PDH, a housekeeping enzyme, providing the NADPH necessary for the synthesis of steroids. The studies were combined with an analysis of testosterone level.

Material and Methods

The experiments involved 20 sexually mature male rats of the inbred Wistar strain. The animals were kept in standard conditions: temperature 22°C, air humidity 50-60%, and the 12/12 hours L/D light regime. The animals were fed with standard fodder and had tap water available *ad libitum*. The rats were divided into two groups of 10 animals in each: control and experimental. The experimental rats received naringenin subcutaneously at a dose of 15 mg/kg b.m. in 0.2 ml dimethylsulphoxide (DMSO) each day for 14 consecutive days. The control rats were given subcutaneous injections of DMSO. After 14 days, the rats were decapitated and their testes removed and frozen in liquid nitrogen. The testes were then cut into 8 μ m sections in a cryostat and subjected to histochemical reactions to detect $\Delta^3\beta$ HSD (EC 1.1.1.145) [5], 17β HSD (EC 1.1.1.51) [6,7], and G6PDH (EC 1.1.1.49) [8]. The enzyme activity in Leydig cells was assessed by the microdensitometric method. A computer-assisted image analyser, with the Multi Scan 6.08 software and an 8-bit grey scale, was used for the assessment. The intensity of the histoenzymatic reaction was estimated by measuring the integrated optical density (IOD) in the marked area. The intensity of histochemical reactions, shown on microphotographs, was assessed as either weak, moderate, or strong.

A portion of the material was placed aside to carry out a radioimmunological assay for testosterone level. This determi-

ADDRESS FOR CORRESPONDENCE:

Papież Monika
Department of Cytobiology and Histochemistry
Pharmaceutical Faculty, Medical College,
Jagiellonian University
Medyczna 9, 30-688 Kraków, Poland
e-mail: mfpapiez@cyf-kr.edu.pl

nation was completed with the use of a standard kit (from DSL) [9]. The significant differences between the groups were statistically tested, using ANOVA procedure.

Results

The greatest intensity of enzymatic reactions occurred in Leydig cells. The activity in the seminiferous epithelium was either weak or absent, depending on the examined enzyme.

The activity of ^{3}HSD in Leydig cells of the control rats was strong (Fig. 1a), whereas in the rats, treated with naringenin, it was either high or moderate. The activity of ^{17}HSD in Leydig cells of the control rats was moderate, while the intensity of that reaction in the experimental rats was weak (Fig. 1 b,c). The activity of G6PDH in Leydig cells of the control rats was either moderate or high, whereas histochemical reaction in the experimental animals was moderate (Fig. 1d,e). Microdensitometric measurements showed a statistically significant decrease in ^{3}HSD and G6PDH activities in the experimental rats, compared with the respective values in the control animals (Fig. 2). The difference between the control and the experimental group in the ^{3}HSD was not significant. In the homogenates of testes of the experimental rats, a statistically insignificant increase in testosterone content (120 ± 45 ng/g tissue) was found, compared with that in the control group (82 ± 26 ng/g tissue).

Discussion

This paper presents preliminary results of a study on the effects of naringenin *in vivo*, on the activity of selected enzymes, involved in the biosynthetic function of rat testes.

The observed drop of ^{17}HSD activity in the rats, receiving naringenin, may indicate a hormone-like effect of this flavonoid. It is known from literature that the administration of $^{17}\text{-estradiol}$ to male rats also leads to a drop in steroidogenic enzymes [10]. The obtained results match those of Le Bail et al. [4], who demonstrated that naringenin inhibited the activity of ^{17}HSD and aromatase in the microsomes, isolated from the human placenta. The application of naringenin to the rats did not result in any changes in the activity of ^{3}HSD , compared to respective values in the control. The results agree with those, reported by Le Bail et al. [11], who demonstrated that naringenin *in vitro* had not affected the activity of ^{3}HSD , an enzyme, involved in steroidogenesis. Despite the drop in ^{17}HSD activity, in the rats receiving naringenin, a slight increase in testosterone level occurred in the testes of those individuals. One may thus presume that, in rats, naringenin inhibits the activity of the enzymes, participating in the metabolism of testosterone, such as aromatase and ^{17}HSD . It should be added that the effect, exerted by phytoestrogens on the synthesis and release of androgens, has not yet been determined and the results of studies often contradict one another [12, 13]. In the rats, receiving naringenin, a decrease in the level of anabolic processes was noted, as indicated by drop in the activity of G6PDH dehydrogenase. The decrease in

Figure 1. (a) Histochemical reaction for ^{3}HSD (200x). Strong reaction in Leydig cells (arrow) of a control rat. (b,c) Histochemical reaction for ^{17}HSD (320x). The arrows show Leydig cells. (b) strong reaction in Leydig cells of a control rat. (c) Moderate and weak reaction in Leydig cells of an experimental rat. (d,e) Histochemical reaction for G6PDH (200x). (d) Moderate reaction in Leydig cells (arrow) of a control rat. (e) Weak reaction in Leydig cells (arrow) of an experimental rat.

Figure 2. The activity of dehydrogenases in Leydig cells of the control and naringenin treated rats. Microdensitometrical analysis. Values are the means \pm SD. Statistically significant differences as compared to the controls. * $p < 0.05$, ** $p < 0.01$.

Acknowledgements

The author is grateful to the staff of the Department of Clinical Biochemistry, Institute of Paediatrics, The Jagiellonian University, for the performed radioimmunological analysis. This study was supported by Grant No. BW/262/P/F/2004 from the State Committee for Scientific Research (KBN).

References

1. Tomas-Barberan FA, Clifford MN. Flavanones, chalcones and dihydrochalcones - nature, occurrence and dietary burden. *J Sci Food and Agric*, 2000; 80: 1073-80.
2. Schaefer O, Humpel M, Fritzemeier KH, Bohlmann R, Schleuning WD. 8-Prenyl naringenin is a potent ER α selective phytoestrogen present in hops and beer. *J Steroid Biochem*, 2003; 84: 359-60.
3. Breinholt VM, Svendsen GW, Dragsted LO, Hossaini A. The citrus-derived flavonoid naringenin exerts uterotrophic effects in female mice at human relevant doses. *Basic Clin Pharmacol Toxicol*, 2004; 94: 30-6.
4. Le Bail JC, Laroche T, Fournier F, Habrioux G. Aromatase and 17 β -hydroxysteroid dehydrogenase inhibited by flavonoids. *Cancer Lett*, 1998; 133: 101-6.
5. Levy H, Wendler-Deane H, Rubin BL. Visualization of steroid-3 β -ol-dehydrogenase activity in tissues of intact and hypophysectomized rats. *Endocrinology*, 1959; 65: 932-43.
6. Erpino MJ. Effects of substrate on histochemistry of 3 β -HSD in testes. *Gen Comp Endocr*, 1971; 17: 563-6.
7. Schafers BA, Schlutius BG, Haider SG. Ontogenesis of oxidative reaction of 17 β -hydroxysteroid dehydrogenase and 11 β -hydroxysteroid dehydrogenase in rat Leydig cells, a histochemical study. *Histochem J*, 2001; 33: 585-95.
8. Pearse AGE. *Histochemistry: Theoretical and Applied*. Vol 2. 3th ed. London: Churchill-Livingstone: 1972.
9. Yalow R, Berson S. Introduction and general considerations. In: Odel WD, Daughday WH, editors. *Principles of competitive protein binding assay*. Philadelphia: JB Lippincott Co.; 1971.
10. Brinkmann AO, Leemberg FG, Roodnat EM, De Joung FH, Van der Molen HJ. A specific action of estradiol on enzymes involved in testicular steroidogenesis. *Biol Reprod*, 1980; 23: 801-9.
11. Le Bail J Ch, Champavier Y, Chulia A-J, Habrioux G. Effects of phytoestrogens on aromatase, 3 β and 17 β -hydroxysteroid dehydrogenase activities and human breast cancer cells. *Life Sci*, 2000; 66: 1281-91.
12. Roberts D, Veeramachaneni DN, Schlaff WD, Awoniyi CA. Effects of chronic dietary exposure to genistein, a phytoestrogen, during various stages of development on reproductive hormones and spermatogenesis in rats. *Endocrine*, 2000; 13: 281-6.
13. Papież M. The influence of hollyhock extract administration on testicular function in rats. *J Mol Histol*, 2004; 35: 1-8 (in press).
14. Vanitha-Kumari G. Effect of para-chlorophenylalanine on male rats: histopathological and biochemical changes in the testes. *Indian J Physiol Pharmacol*, 1986; 30: 223-31.